

# Geometric control and applications

Ludovic Rifford

Université Nice Sophia Antipolis

Fields Institute - November 6-7, 2014

- Lecture 1:

A controllability result: The Chow-Rashevsky Theorem


- Lecture 2:

An optimal control study: Sub-Riemannian geodesics

# Lecture 1

## A controllability result: The Chow-Rashevsky Theorem

# Control of an inverted pendulum



# Control systems

A general control system has the form

$$\dot{x} = f(x, u)$$

where

- $x$  is the state in  $M$
- $u$  is the control in  $U$

## Proposition

*Under classical assumptions on the data, for every  $x \in M$  and every measurable control  $u : [0, T] \rightarrow U$  the Cauchy problem*

$$\begin{cases} \dot{x}(t) = f(x(t), u(t)) & \text{a.e. } t \in [0, T], \\ x(0) = x \end{cases}$$

*admits a unique solution*

$$x(\cdot) = x(\cdot; x, u) : [0, T] \longrightarrow M.$$


# Controllability issues

Given two points  $x_1, x_2$  in the state space  $M$  and  $T > 0$ , can we find a control  $u$  such that the solution of

$$\begin{cases} \dot{x}(t) = f(x(t), u(t)) & \text{a.e. } t \in [0, T] \\ x(0) = x_1 \end{cases}$$

satisfies

$$x(T) = x_2 \quad ?$$



# Controllability of linear control systems in $\mathbb{R}^n$

An autonomous linear control system in  $\mathbb{R}^n$  has the form

$$\dot{\xi} = A\xi + B u,$$

with  $\xi \in \mathbb{R}^n$ ,  $u \in \mathbb{R}^m$ ,  $A \in M_n(\mathbb{R})$ ,  $B \in M_{n,m}(\mathbb{R})$ .

## Theorem

*The following assertions are equivalent:*

- (i) *For any  $T > 0$  and any  $\xi_1, \xi_2 \in \mathbb{R}^n$ , there is  $u \in L^1([0, T]; \mathbb{R}^m)$  such that*

$$\xi(T; \xi_1, u) = \xi_2.$$

- (ii) *The Kalman rank condition is satisfied:*

$$\text{rk}(B, AB, A^2B, \dots, A^{n-1}B) = n.$$

# Proof of the theorem

## Duhamel's formula

$$\xi(T; \xi, \mathbf{u}) = e^{TA} \xi + e^{TA} \int_0^T e^{-tA} B \mathbf{u}(t) dt.$$

Then the controllability property (i) is equivalent to the surjectivity of the mappings

$$\mathcal{F}^T : \mathbf{u} \in L^1([0, T]; \mathbb{R}^m) \mapsto \int_0^T e^{-tA} B \mathbf{u}(t) dt.$$

# Proof of (ii) $\Rightarrow$ (i)

If  $\mathcal{F}^T$  is not onto (for some  $T > 0$ ), there is  $p \neq 0_n$  such that

$$\left\langle p, \int_0^T e^{-tA} B u(t) dt \right\rangle = 0 \quad \forall u \in L^1([0, T]; \mathbb{R}^m).$$

Using the linearity of  $\langle \cdot, \cdot \rangle$  and taking  $u(t) = B^* e^{-tA} p$ , we infer that

$$p^* e^{-tA} B = 0 \quad \forall t \in [0, T].$$

Derivating  $n$  times at  $t = 0$  yields

$$p^* B = p^* A B = p^* A^2 B = \dots = p^* A^{n-1} B = 0.$$

Which means that  $p$  is orthogonal to the image of the  $n \times mn$  matrix

$$(B, AB, A^2 B, \dots, A^{n-1} B).$$

**Contradiction !!!**

# Proof of (i) $\Rightarrow$ (ii)

If

$$\text{rk}(B, AB, A^2B, \dots, A^{n-1}B) < n,$$

there is a nonzero vector  $p$  such that

$$p^* B = p^* AB = p^* A^2B = \dots = p^* A^{n-1}B = 0.$$

By the Cayley-Hamilton Theorem, we deduce that

$$p^* A^k B = 0 \quad \forall k \geq 1,$$

and in turn

$$p^* e^{-tA} B = 0 \quad \forall t \geq 0.$$

We infer that

$$\left\langle p, \int_0^T e^{-tA} B u(t) dt \right\rangle = 0 \quad \forall u \in L^1([0, T]; \mathbb{R}^m), \forall T > 0.$$

**Contradiction !!!**

# Application to local controllability

Let  $\dot{x} = f(x, u)$  be a nonlinear control system with  $x \in \mathbb{R}^n$ ,  $u \in \mathbb{R}^m$  and  $f : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^n$  of class  $C^1$ .

## Theorem


Assume that  $f(x_0, 0) = 0$  and that the pair

$$A = \frac{\partial f}{\partial x}(x_0, 0), \quad B = \frac{\partial f}{\partial u}(x_0, 0),$$

satisfies the Kalman rank condition. Then for there is  $\delta > 0$  such that for any  $x_1, x_2$  with  $|x_1 - x_0|, |x_2 - x_0| < \delta$ , there is  $u : [0, 1] \rightarrow \mathbb{R}^m$  smooth satisfying

$$x(1; x_1, u) = x_2.$$

# Local controllability around $x_0$



# Proof of the Theorem

Define  $\mathcal{G} : \mathbb{R}^n \times L^1([0, 1]; \mathbb{R}^m) \rightarrow \mathbb{R}^n \times \mathbb{R}^n$  by


$$\mathcal{G}(x, u) := (x, x(1; x, u)).$$

The mapping  $\mathcal{G}$  is a  $C^1$  submersion at  $(0, 0)$ . Thus there are  $n$  controls  $u^1, \dots, u^n$  in  $L^1([0, 1]; \mathbb{R}^m)$  such that

$$\begin{aligned} \tilde{\mathcal{G}} : \mathbb{R}^n \times \mathbb{R}^n &\longrightarrow \mathbb{R}^n \times \mathbb{R}^n \\ (x, \lambda) &\longmapsto \mathcal{G}(x, \sum_{i=1}^n \lambda_i u^i) \end{aligned}$$

is a  $C^1$  diffeomorphism at  $(0, 0)$ . Since the set of smooth controls is dense in  $L^1([0, 1]; \mathbb{R}^m)$ , we can take  $u^1, \dots, u^n$  to be smooth. **We apply the Inverse Function Theorem.**

# Back to the inverted pendulum



The equations of motion are given by

$$\begin{aligned}(M + m)\ddot{x} + ml\ddot{\theta}\cos\theta - ml\dot{\theta}^2\sin\theta &= u \\ ml^2\ddot{\theta} - mgl\sin\theta + ml\ddot{x}\cos\theta &= 0.\end{aligned}$$

# Back to the inverted pendulum

The linearized control system at  $x = \dot{x} = \theta = \dot{\theta} = 0$  is given by

$$\begin{aligned}(M + m)\ddot{x} + m\ell\ddot{\theta} &= u \\ m\ell^2\ddot{\theta} - mg\ell\theta + m\ell\ddot{x} &= 0.\end{aligned}$$

It can be written as a control system

$$\dot{\xi} = A\xi + B u,$$

with  $\xi = (x, \dot{x}, \theta, \dot{\theta})$ ,

$$A = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & -\frac{mg}{M} & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & \frac{(M+m)g}{M\ell} & 0 \end{pmatrix} \quad \text{and} \quad B = \begin{pmatrix} 0 \\ \frac{1}{M} \\ 0 \\ -\frac{1}{M\ell} \end{pmatrix}.$$

# Back to the inverted pendulum

The Kalman matrix  $(B, AB, A^2, A^3B)$  equals

$$\begin{pmatrix} 0 & \frac{1}{M} & 0 & \frac{mg}{M^2\ell} \\ \frac{1}{M} & 0 & \frac{mg}{M^2\ell} & 0 \\ 0 & -\frac{1}{M\ell} & 0 & -\frac{(M+m)g}{M^2\ell^2} \\ -\frac{1}{M\ell} & 0 & -\frac{(M+m)g}{M^2\ell^2} & 0 \end{pmatrix}.$$

Its determinant equals

$$-\frac{g^2}{M^4\ell^4} < 0$$

In conclusion, the inverted pendulum is locally controllable around  $(0, 0, 0, 0)^*$ .

# Movie

# The Chow-Rashevsky Theorem

## Theorem (Chow 1939, Rashevsky 1938)

Let  $M$  be a smooth manifold and  $X^1, \dots, X^m$  be  $m$  smooth vector fields on  $M$ . Assume that

$$\text{Lie} \{X^1, \dots, X^m\} (x) = T_x M \quad \forall x \in M.$$


Then the control system

$$\dot{x} = \sum_{i=1}^m u_i X^i(x)$$

is *locally controllable in any time at every point* of  $M$ .

# Comment 1

The **local controllability in any time at every point** means that for every  $x_0 \in M$ , every  $T > 0$  and every neighborhood  $U$  of  $x_0$ , there is a neighborhood  $V \subset U$  of  $x_0$  such that for any  $x_1, x_2 \in V$ , there is a control  $u \in L^1([0, T]; \mathbb{R}^m)$  such that the trajectory  $x(\cdot; x_1, u) : [0, T] \rightarrow M$  remains in  $U$  and steers  $x_1$  to  $x_2$ , i.e.  $x(T; x_1, u) = x_2$ .



**Local controllability in time  $T > 0$**

$\Rightarrow$  **Local controllability in time  $T' > 0$ ,  $\forall T' > 0$**

# Comment II

If  $M$  is **connected** then

**Local controllability  $\Rightarrow$  Global controllability**

Let  $x \in M$  be fixed. Denote by  $\mathcal{A}(x)$  the accessible set from  $x$ , that is

$$\begin{aligned}\mathcal{A}(x) &:= \{x(T; x, u) \mid T \geq 0, u \in L^1\} \\ &= \{x(1; x, u) \mid u \in L^1\}.\end{aligned}$$

- By local controllability,  $\mathcal{A}(x)$  is open.
- Let  $y$  be in the closure of  $\mathcal{A}(x)$ . The set  $\mathcal{A}(y)$  contains a small ball centered at  $y$  and there are points of  $\mathcal{A}(x)$  in that ball. Then  $\mathcal{A}(x)$  is closed.

**By connectedness of  $M$ , we infer that  $\mathcal{A}(x) = M$  for every  $x \in M$ , and in turn that the control system is globally controllable in any time.**

# The Chow-Rashevsky Theorem

Theorem (Chow 1939, Rashevsky 1938)

Let  $M$  be a smooth manifold and  $X^1, \dots, X^m$  be  $m$  smooth vector fields on  $M$ . Assume that

$$\text{Lie}\{X^1, \dots, X^m\}(x) = T_x M \quad \forall x \in M.$$

Then the control system  $\dot{x} = \sum_{i=1}^m u_i X^i(x)$  is locally controllable in any time at every point of  $M$ .

The condition in red is called Hörmander's condition or bracket generating condition. Families of vector fields satisfying that condition are called nonholonomic, completely nonholonomic, or totally nonholonomic.

## Definition

Given two smooth vector fields  $X, Y$  on  $\mathbb{R}^n$ , the Lie bracket  $[X, Y]$  at  $x \in \mathbb{R}^n$  is defined by


$$[X, Y](x) = DY(x)X(x) - DX(x)Y(x).$$

The Lie brackets of two smooth vector fields on  $M$  can be defined in charts with the above formula.


Given a family  $\mathcal{F}$  of smooth vector fields on  $M$ , we denote by  $\text{Lie}\{\mathcal{F}\}$  the Lie algebra generated by  $\mathcal{F}$ . It is the smallest vector subspace  $S$  of smooth vector fields containing  $\mathcal{F}$  that also satisfies

$$[X, Y] \in S \quad \forall X \in \mathcal{F}, \forall Y \in S.$$


# Comment III




# Comment III




# Comment III



# Comment III



# Comment III





# Comment III

Given a family  $\mathcal{F}$  of smooth vector fields on  $M$ , we set  $\text{Lie}^1(\mathcal{F}) := \text{Span}(\mathcal{F})$ , and define recursively  $\text{Lie}^k(\mathcal{F})$  ( $k = 2, 3, \dots$ ) by

$$\text{Lie}^{k+1}(\mathcal{F}) := \text{Span}\left(\text{Lie}^k(\mathcal{F}) \cup \left\{ [X, Y] \mid X \in \mathcal{F}, Y \in \text{Lie}^k(\mathcal{F}) \right\}\right).$$

We have

$$\text{Lie}\{\mathcal{F}\} = \bigcup_{k \geq 1} \text{Lie}^k(\mathcal{F}).$$

For example, the Lie algebra  $\text{Lie}\{X^1, \dots, X^m\}$  is the vector subspace of smooth vector fields which is spanned by all the brackets (made from  $X^1, \dots, X^m$ ) of length  $1, 2, 3, \dots$

Since  $M$  has finite dimension, for every  $x \in M$ , there is  $r = r(x) \geq 1$  (called degree of nonholonomy at  $x$ ) such that

$$T_x M \supset \text{Lie}\{X^1, \dots, X^m\}(x) = \text{Lie}^r\{X^1, \dots, X^m\}(x).$$

## Comment IV

We can prove the Chow-Rashevsky Theorem in the contact case in  $\mathbb{R}^3$  as follows:

### Exercise

Let  $X^1, X^2$  be two smooth vector fields in  $\mathbb{R}^3$  such that

$$\text{Span}\{X^1(0), X^2(0), [X^1, X^2](0)\} = \mathbb{R}^3.$$

Then the mapping  $\varphi_\lambda : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  defined by

$$\varphi_\lambda(t_1, t_2, t_3) = e^{\lambda X^1} \circ e^{t_3 X^2} \circ e^{-\lambda X^1} \circ e^{t_2 X^2} \circ e^{t_1 X^1}(0)$$

is a local diffeomorphism at the origin for  $\lambda > 0$  small.

$\rightsquigarrow$  Ball-Box Theorem

# The End-Point mapping

Given a control system of the form

$$\dot{x} = \sum_{i=1}^m u_i X^i(x) \quad (x \in M, u \in \mathbb{R}^m),$$

we define the **End-Point mapping** from  $x$  in time  $T > 0$  as

$$\begin{aligned} E^{x,T} : L^2([0, T]; \mathbb{R}^m) &\longrightarrow M \\ u &\longmapsto x(T; x, u) \end{aligned}$$

## Proposition

*The mapping  $E^{x,T}$  is of class  $C^1$  (on its domain) and*

$$D_u E^{x,T}(v) = \xi(T), \quad \text{where}$$

$$\dot{\xi} = \left( \sum_{i=1}^m u_i D_{x_u} X^i \right) \cdot \xi + \sum_{i=1}^m v_i X^i(x_u), \quad \xi(0) = 0.$$

# Linearized control system

## Remark

Setting for every  $t \in [0, T]$ ,  $A_u(t) := \sum_{i=1}^m u_i(t) D_{x_u(t)} X^i$ , we have

$$D_u E^{x,T}(v) = S_u(T) \int_0^T S_u(t)^{-1} \sum_{i=1}^m v_i(t) X^i(x_u(t)) dt$$

with  $S_u$  solution of  $\dot{S}_u = A_u S_u$  a.e.  $t \in [0, T]$ ,  $S_u(0) = I_n$ .

## Proposition

For every  $u \in L^2([0, T]; \mathbb{R}^m)$  and any  $i = 1, \dots, m$ , we have

$$X^i(E^{x,T}(u)) \in D_u E^{x,T}(L^2([0, T]; \mathbb{R}^m)).$$


# Regular controls vs. Singular controls

## Definition

A control  $u \in L^2([0, T]; \mathbb{R}^m)$  is called **regular** with respect to  $E^{x, T}$  if  $E^{x, T}$  is a submersion at  $u$ . If not,  $u$  is called **singular**.

## Exercise

*The concatenations  $u^1 * u^2$  and  $u^2 * u^1$  of a regular control  $u^1$  with another control  $u^2$  are regular.*



# Rank of a control


## Definition

The rank of a control  $u \in L^2([0, T]; \mathbb{R}^m)$  (with respect to  $E^{x, T}$ ) is defined as the dimension of the image of the linear mapping  $D_u E^{x, T}$ . We denote it by  $\text{rank}^{x, T}(u)$ .

## Exercise

The following properties hold:

- $\text{rank}^{x, T_1+T_2}(u^1 * u^2) \geq \max\{\text{rank}^{x, T_1}(u^1), \text{rank}^{y, T_2}(u^2)\}$ .
- $\text{rank}^{y, T_1}(\check{u}^1) = \text{rank}^{x, T_1}(u^1)$ .



# Openness: Statement

The Chow-Rashevsky will follow from the following result:

## Proposition

Let  $M$  be a smooth manifold and  $X^1, \dots, X^m$  be  $m$  smooth vector fields on  $M$ . Assume that

$$\text{Lie} \{X^1, \dots, X^m\} (x) = T_x M \quad \forall x \in M.$$

Then, for every  $x \in M$  and every  $T > 0$ , the End-Point mapping

$$\begin{aligned} E^{x,T} : L^2([0, T]; \mathbb{R}^m) &\longrightarrow M \\ u &\longmapsto x(T; x, u) \end{aligned}$$

is open (on its domain).

# Openness: Sketch of proof

Let  $x \in M$  and  $T > 0$  be fixed. Set for every  $\epsilon > 0$ ,

$$d(\epsilon) = \max \left\{ \text{rank}^{x, \epsilon}(u) \mid \|u\|_{L^2} < \epsilon \right\}.$$

**Claim:**  $d(\epsilon) = n \quad \forall \epsilon > 0$ .

If not, we have  $d(\epsilon) = d_0 \in \{1, \dots, n-1\}$  for some  $\epsilon > 0$ .

Given  $u^\epsilon$  s.t.  $\text{rank}^{x, \epsilon}(u^\epsilon) = d_0$ , there are  $d_0$  controls  $v^1, \dots, v^{d_0}$  such that the mapping

$$\mathcal{E} : \lambda = (\lambda^1, \dots, \lambda^{d_0}) \in \mathbb{R}^{d_0} \mapsto E^{x, \epsilon} \left( u^\epsilon + \sum_{j=1}^{d_0} \lambda^j v^j \right)$$


is an immersion near 0. Thus, its local image  $N$  is a  $d_0$  dimensional submanifold of  $M$  of class  $C^1$  such that

$$X^i(\mathcal{E}(\lambda)) \in \text{Im}(D_\lambda \mathcal{E}) = T_y N. \quad \text{Contradiction!!!}$$

# Openness: Sketch of proof (the return method)

To conclude, we pick (for any  $\epsilon > 0$  small) a regular control  $u^\epsilon$  in  $L^2([0, \epsilon]; \mathbb{R}^m)$  and define  $\tilde{u} \in L^2([0, T + 2\epsilon]; \mathbb{R}^m)$  by

$$\tilde{u} := u^\epsilon * \check{u}^\epsilon * u.$$



Up to reparametrizing  $u$  into a control  $v$  on  $[0, T - 2\epsilon]$ , the new control  $\tilde{v} = u^\epsilon * \check{u}^\epsilon * v$  is regular, close to  $u$  in  $L^2$  provided  $\epsilon > 0$  is small, and steers  $x$  to  $E^{x,T}(u)$ .

The openness follows from the Inverse Function Theorem.

## Proposition

Let  $M$  be a smooth manifold and  $X^1, \dots, X^m$  be  $m$  smooth vector fields on  $M$ . Assume that

$$\text{Lie} \{X^1, \dots, X^m\} (x) = T_x M \quad \forall x \in M.$$

Then, for every  $x \in M$  and every  $T > 0$ , the set of controls which are regular w.r.t.  $E^{x,T}$  is open and dense in  $L^2$ .

The above result holds indeed in the smooth topology.

## Proposition (Sontag)

Under the same assumptions, the set of controls which are regular w.r.t.  $E^{x,T}$  is open and dense in  $C^\infty$ .

# Example: The baby stroller




$$\begin{cases} \dot{x} = u_1 \cos \theta \\ \dot{y} = u_1 \sin \theta \\ \dot{\theta} = u_2 \end{cases}$$


$$X = \begin{pmatrix} \cos \theta \\ \sin \theta \\ 0 \end{pmatrix}, \quad Y = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, \quad [X, Y] = \begin{pmatrix} -\sin \theta \\ \cos \theta \\ 0 \end{pmatrix}$$

$$\text{Span}\{X(\xi), Y(\xi), [X, Y](\xi)\} = \mathbb{R}^3 \quad \forall \xi = (x, y, \theta).$$

# Example: The baby stroller



# Example: The baby stroller



Thank you for your attention !!

## Lecture 2

# Sub-Riemannian geodesics

# Sub-Riemannian structures

Let  $M$  be a smooth connected manifold of dimension  $n \geq 2$ .

## Definition

A sub-Riemannian structure on  $M$  is a pair  $(\Delta, g)$  where:

- $\Delta$  is a **totally nonholonomic distribution** of rank  $m \in [2, n]$ , that is it is defined locally as

$$\Delta(x) = \text{Span}\{X^1(x), \dots, X^m(x)\} \subset T_x M,$$

where  $X^1, \dots, X^m$  are  $m$  linearly independent vector fields satisfying the Hörmander condition.

- $g_x$  is a **scalar product** on  $\Delta(x)$ .

# Sub-Riemannian structures

## Remark

- In general  $\Delta$  does not admit a global frame. However we can always construct  $k = m \cdot (n + 1)$  smooth vector fields  $Y^1, \dots, Y^k$  such that

$$\Delta(x) = \text{Span}\{Y^1(x), \dots, Y^k(x)\} \quad \forall x \in M.$$

- If  $(M, g)$  is a Riemannian manifold, then any totally nonholomic distribution  $\Delta$  gives rise to a SR structure  $(\Delta, g)$  on  $M$ .

## Example (Heisenberg)

Take in  $\mathbb{R}^3$ ,  $\Delta = \text{Span}\{X^1, X^2\}$  with

$$X^1 = \partial_x - \frac{y}{2}\partial_z, \quad X^2 = \partial_y + \frac{x}{2}\partial_z \quad \text{and} \quad g = dx^2 + dy^2.$$

# The Chow-Rashevsky Theorem

## Definition

We call **horizontal path** any path  $\gamma \in W^{1,2}([0, 1]; M)$  satisfying

$$\dot{\gamma}(t) \in \Delta(\gamma(t)) \quad \text{a.e. } t \in [0, 1].$$

We observe that if  $\Delta = \text{Span}\{Y^1, \dots, Y^k\}$ , for any  $x \in M$  and any control  $u \in L^2([0, 1]; \mathbb{R}^k)$ , the solution to

$$\dot{\gamma} = \sum_{i=1}^k u_i Y^i(\gamma), \quad \gamma(0) = x$$

is an horizontal path joining  $x$  to  $\gamma(1)$ .

## Theorem (Chow-Rashevsky)

*Let  $\Delta$  be a totally nonholonomic distribution on  $M$  then any pair of points can be joined by an horizontal path.*

# The sub-Riemannian distance

The **length** (w.r.t  $g$ ) of an horizontal path  $\gamma$  is defined as

$$\text{length}^g(\gamma) := \int_0^T |\dot{\gamma}(t)|_{\gamma(t)}^g dt$$

## Definition

Given  $x, y \in M$ , the **sub-Riemannian distance** between  $x$  and  $y$  is

$$d_{SR}(x, y) := \inf \left\{ \text{length}^g(\gamma) \mid \gamma \text{ hor.}, \gamma(0) = x, \gamma(1) = y \right\}.$$

## Proposition

*The manifold  $M$  equipped with the distance  $d_{SR}$  is a metric space whose topology coincides with the topology of  $M$  (as a manifold).*

# Minimizing horizontal paths and geodesics

## Definition

Given  $x, y \in M$ , we call **minimizing horizontal path** between  $x$  and  $y$  any horizontal path  $\gamma : [0, T] \rightarrow M$  connecting  $x$  to  $y$  such that

$$d_{SR}(x, y) = \text{length}^g(\gamma).$$

The **sub-Riemannian energy** between  $x$  and  $y$  is defined as

$$e_{SR}(x, y) := \inf \left\{ \text{energy}^g(\gamma) := \int_0^1 \left( |\dot{\gamma}(t)|_{g_{\gamma(t)}}^g \right)^2 dt \mid \gamma \dots \right\}.$$

## Definition

We call **minimizing geodesic** between  $x$  and  $y$  any horizontal path  $\gamma : [0, 1] \rightarrow M$  connecting  $x$  to  $y$  such that

$$e_{SR}(x, y) = \text{energy}^g(\gamma).$$

# A SR Hopf-Rinow Theorem

## Theorem

Let  $(\Delta, g)$  be a sub-Riemannian structure on  $M$ . Assume that  $(M, d_{SR})$  is a complete metric space. Then the following properties hold:

- The closed balls  $\bar{B}_{SR}(x, r)$  are compact (for any  $r \geq 0$ ).
- For every  $x, y \in M$ , there exists at least one minimizing geodesic joining  $x$  to  $y$ .


## Remark

Given a complete Riemannian manifold  $(M, g)$ , for any totally nonholonomic distribution  $\Delta$  on  $M$ , the SR structure  $(\Delta, g)$  is complete. *As a matter of fact, since  $d_g \leq d_{SR}$  any Cauchy sequence w.r.t.  $d_{SR}$  is Cauchy w.r.t.  $d_g$ .*

# The Hamiltonian geodesic equation

Let  $x, y \in M$  and a **minimizing geodesic**  $\bar{\gamma}$  joining  $x$  to  $y$  be fixed. The SR structure admits **an orthonormal frame** along  $\bar{\gamma}$ , that is there is an open neighborhood  $\mathcal{V}$  of  $\bar{\gamma}([0, 1])$  and an orthonormal family of  $m$  vector fields  $X^1, \dots, X^m$  such that

$$\Delta(z) = \text{Span}\{X^1(z), \dots, X^m(z)\} \quad \forall z \in \mathcal{V}.$$



# The Hamiltonian geodesic equation

There is a control  $\bar{u} \in L^2([0, 1]; \mathbb{R}^m)$  such that

$$\dot{\bar{\gamma}}(t) = \sum_{i=1}^m \bar{u}_i(t) X^i(\bar{\gamma}(t)) \quad \text{a.e. } t \in [0, 1].$$

Moreover, on the one hand any control  $u \in \mathcal{U} \subset L^2([0, 1]; \mathbb{R}^m)$  ( $u$  sufficiently close to  $\bar{u}$ ) gives rise to a trajectory  $\gamma_u$  solution of

$$\dot{\gamma}_u = \sum_{i=1}^m u^i X^i(\gamma_u) \quad \text{on } [0, T], \quad \gamma_u(0) = x.$$

On the other hand, for any horizontal path  $\gamma : [0, 1] \rightarrow \mathcal{V}$  there is a (unique) control  $u \in L^2([0, 1]; \mathbb{R}^m)$  for which the equation in red is satisfied.

# The Hamiltonian geodesic equation

So, considering as previously the **End-Point mapping**

$$E^{x,1} : L^2([0, 1]; \mathbb{R}^m) \longrightarrow M$$

defined by

$$E^{x,1}(u) := \gamma_u(1),$$

and setting  $C(u) = \|u\|_{L^2}^2$ , we observe that  $\bar{u}$  is solution to the following **optimization problem with constraints**:

$\bar{u}$  minimizes  $C(u)$  among all  $u \in \mathcal{U}$  s.t.  $E^{x,1}(u) = y$ .

(Since the family  $X^1, \dots, X^m$  is orthonormal, we have

$$\text{energy}^g(\gamma_u) = C(u) \quad \forall u \in \mathcal{U}.)$$

# The Hamiltonian geodesic equation

## Proposition (Lagrange Multipliers)

There are  $p \in T_y^*M \simeq (\mathbb{R}^n)^*$  and  $\lambda_0 \in \{0, 1\}$  with  $(\lambda_0, p) \neq (0, 0)$  such that

$$p \cdot D_{\bar{u}} E^{x,1} = \lambda_0 D_{\bar{u}} C.$$

## Proof.

The mapping  $\Phi : \mathcal{U} \rightarrow \mathbb{R} \times M$  defined by

$$\Phi(u) := (C(u), E^{x,1}(u))$$

cannot be a submersion at  $\bar{u}$ . As a matter of fact, if  $D_{\bar{u}}\Phi$  is surjective, then it is open at  $\bar{u}$ , so it must contain elements of the form  $(C(\bar{u}) - \delta, y)$  for  $\delta > 0$  small.  $\square$

$\rightsquigarrow$  two cases:  $\lambda_0 = 0$  or  $\lambda_0 = 1$ .

# The Hamiltonian geodesic equation

**First case:**  $\lambda_0 = 0$

Then we have

$$p \cdot D_{\bar{u}} E^{x,1} = 0 \text{ with } p \neq 0.$$

So  $\bar{u}$  is **singular** (w.r.t.  $x$  and  $T = 1$ ).

## Remark

- *If  $\Delta$  has rank  $n$ , that is  $\Delta = TM$  (Riemannian case), then there are no singular control. So this case cannot occur.*
- *If there are no nontrivial singular control, then this case cannot occur.*
- *If there are no nontrivial singular minimizing control, then this case cannot occur.*

# The Hamiltonian geodesic equation

**Second case:**  $\lambda_0 = 1$

Define the Hamiltonian  $H : \mathcal{V} \times (\mathbb{R}^n)^* \rightarrow \mathbb{R}$  by

$$H(x, p) := \frac{1}{2} \sum_{i=1}^m (p \cdot X^i(x))^2.$$

## Proposition

There is a smooth arc  $p : [0, 1] \rightarrow (\mathbb{R}^n)^*$  with  $p(1) = p/2$  such that

$$\begin{cases} \dot{\bar{\gamma}} &= \frac{\partial H}{\partial p}(\bar{\gamma}, p) = \sum_{i=1}^m [p \cdot X^i(\bar{\gamma})] X^i(\bar{\gamma}) \\ \dot{p} &= -\frac{\partial H}{\partial x}(\bar{\gamma}, p) = -\sum_{i=1}^m [p \cdot X^i(\bar{\gamma})] p \cdot D\bar{\gamma} X^i \end{cases}$$

for a.e.  $t \in [0, 1]$  and  $\bar{u}_i(t) = p \cdot X^i(\bar{\gamma}(t))$  for a.e.  $t \in [0, 1]$  and any  $i$ . *In particular, the path  $\bar{\gamma}$  is smooth on  $[0, 1]$ .*

# The Hamiltonian geodesic equation

Proof.

We have  $D_{\bar{u}}C(v) = 2\langle \bar{u}, v \rangle_{L^2}$  and we remember that

$$D_{\bar{u}}E^{x,T}(v) = S(1) \int_0^1 S(t)^{-1} B(t) v(t) dt$$

with

$$\begin{cases} A(t) &= \sum_{i=1}^m u_i(t) D_{\bar{\gamma}(t)} X^i, \\ B(t) &= (X^1(\bar{\gamma}(t)), \dots, X^m(\bar{\gamma}(t))) \end{cases} \quad \forall t \in [0, 1],$$

and  $S$  solution of

$$\dot{S}(t) = A(t)S(t) \text{ for a.e. } t \in [0, 1], \quad S(0) = I_n.$$



# The Hamiltonian geodesic equation

Proof.

Then  $p \cdot D_{\bar{u}} E^{x,1} = \lambda_0 D_{\bar{u}} C$  yields

$$\int_0^1 [p \cdot S(1)S(t)^{-1}B(t) - 2\bar{u}(t)^*] v(t) dt = 0 \quad \forall v \in L^2.$$

We infer that

$$\bar{u}(t) = \frac{1}{2} (p \cdot S(1)S(t)^{-1}B(t))^* \quad \text{a.e. } t \in [0, 1],$$

and that the absolutely continuous arc  $p : [0, 1] \rightarrow (\mathbb{R}^n)^*$  defined by

$$p(t) := \frac{1}{2} p \cdot S(1)S(t)^{-1}$$

satisfies the desired equations. □

# The Hamiltonian geodesic equation

Define the **Hamiltonian**  $H : T^*M \rightarrow \mathbb{R}$  by

$$H(x, p) = \frac{1}{2} \max \left\{ \frac{p(v)^2}{g_x(v, v)} \mid v \in \Delta_x \setminus \{0\} \right\}.$$

We call **normal extremal** any curve  $\psi : [0, T] \rightarrow T^*M$  satisfying

$$\dot{\psi}(t) = \vec{H}(\psi(t)) \quad \forall t \in [0, T].$$

## Theorem

Let  $\gamma : [0, 1] \rightarrow M$  be a minimizing geodesic. One of the two following non-exclusive cases occur:

- $\gamma$  is singular.
- $\gamma$  admits a normal extremal lift.

## Example 1: The Riemannian case

Let  $\Delta(x) = T_x M$  for any  $x \in M$  so that ANY curve is horizontal. There are no singular curve, so any minimizing geodesic is the projection of a normal extremal.

## Example 2: Heisenberg

Recall that in  $\mathbb{R}^3$ ,  $\Delta = \text{Span}\{X^1, X^2\}$  with

$$X^1 = \partial_x - \frac{y}{2}\partial_z, \quad X^2 = \partial_y + \frac{x}{2}\partial_z \text{ and } g = dx^2 + dy^2.$$

# Examples

Any horizontal path has the form  $\gamma_u = (x, y, z) : [0, 1] \rightarrow \mathbb{R}^3$  with

$$\begin{cases} \dot{x}(t) = u_1(t) \\ \dot{y}(t) = u_2(t) \\ \dot{z}(t) = \frac{1}{2} (u_2(t)x(t) - u_1(t)y(t)), \end{cases}$$

for some  $u \in L^2$ . This means that

$$z(1) - z(0) = \int_{\alpha} \frac{1}{2} (x dy - y dx),$$

where  $\alpha$  is the projection of  $\gamma$  to the plane  $z = 0$ . By Stokes' Theorem, we get

$$z(1) - z(0) = \int_{\mathcal{D}} dx \wedge dy + \int_c \frac{1}{2} (x dy - y dx)$$

where  $\mathcal{D}$  is the domain enclosed by  $\alpha$  and the segment  $c = [\alpha(0), \alpha(1)]$ .  $\rightsquigarrow$  **Projections of minimizing horizontal paths must be circles.**

# Examples

Let  $\gamma_u = (x, y, z) : [0, 1] \rightarrow \mathbb{R}^3$  be a minimizing geodesic from  $P_1 := \gamma_u(0)$  to  $P_2 := \gamma_u(1) \neq P_1$ . Since  $u$  is necessarily regular, there is a smooth arc  $p : [0, 1] \rightarrow (\mathbb{R}^3)^*$  s.t.

$$\begin{cases} \dot{x} &= p_x - \frac{v}{2} p_z \\ \dot{y} &= p_y + \frac{x}{2} p_z \\ \dot{z} &= \frac{1}{2} \left( (p_y + \frac{x}{2} p_z) x - (p_x - \frac{v}{2} p_z) y \right), \end{cases} \quad \begin{cases} \dot{p}_x &= - \left( p_y + \frac{x}{2} p_z \right) \frac{p_z}{2} \\ \dot{p}_y &= \left( p_x - \frac{v}{2} p_z \right) \frac{p_z}{2} \\ \dot{p}_z &= 0. \end{cases}$$

Hence  $p_z = \bar{p}_z$  for every  $t$ . Which implies that

$$\ddot{x} = -\bar{p}_z \dot{y} \quad \text{and} \quad \ddot{y} = \bar{p}_z \dot{x}.$$

If  $\bar{p}_z = 0$ , then the geodesic from  $P_1$  to  $P_2$  is a segment with constant speed. If  $\bar{p}_z \neq 0$ , we have or

$$\ddot{x} = -\bar{p}_z^2 \dot{x} \quad \text{and} \quad \ddot{y} = -\bar{p}_z^2 \dot{y}.$$

Which means that the curve  $t \mapsto (x(t), y(t))$  is a circle.

# Examples

## Example 3: The Martinet distribution

In  $\mathbb{R}^3$ , let  $\Delta = \text{Span}\{X^1, X^2\}$  with  $X^1, X^2$  of the form

$$X^1 = \partial_{x_1} \quad \text{and} \quad X^2 = (1 + x_1\phi(x)) \partial_{x_2} + x_1^2 \partial_{x_3},$$

where  $\phi$  is a smooth function and  $g$  be a smooth metric on  $\Delta$ .

### Theorem

*There is  $\bar{\epsilon} > 0$  such that for every  $\epsilon \in (0, \bar{\epsilon})$ , the (singular) horizontal path given by*

$$\gamma(t) = (0, t, 0) \quad \forall t \in [0, \epsilon],$$

*minimizes the length (w.r.t.  $g$ ) among all horizontal paths joining 0 to  $(0, \epsilon, 0)$ . Moreover if  $\{X^1, X^2\}$  is orthonormal w.r.t.  $g$  and  $\phi(0) \neq 0$ , then  $\gamma$  can not be the projection of a normal extremal.*

# The SR exponential mapping

Denote by  $\psi_{x,p} : [0, 1] \rightarrow T^*M$  the solution of

$$\dot{\psi}(t) = \vec{H}(\psi(t)) \quad \forall t \in [0, 1], \quad \psi(0) = (x, p)$$

and let

$$\mathcal{E}_x := \left\{ p \in T_x^*M \mid \psi_{x,p} \text{ defined on } [0, 1] \right\}.$$

## Definition

The **sub-Riemannian exponential map** from  $x \in M$  is defined by

$$\begin{aligned} \exp_x : \mathcal{E}_x \subset T_x^*M &\longrightarrow M \\ p &\longmapsto \pi(\psi_{x,p}(1)). \end{aligned}$$

## Proposition

*Assume that  $(M, d_{SR})$  is complete. Then for every  $x \in M$ ,  $\mathcal{E}_x = T_x^*M$ .*

# On the image of the exponential mapping

## Proposition (Agrachev-Trélat-LR)

Assume that  $(M, d_{SR})$  is complete. Then for every  $x \in M$ , the set  $\exp_x(T_x^*M)$  is open and dense.

## Lemma

Let  $y \neq x$  in  $M$  be such that there is a function  $\phi : M \rightarrow \mathbb{R}$  differentiable at  $y$  such that

$$\phi(y) = d_{SR}^2(x, y) \quad \text{and} \quad d_{SR}^2(x, z) \geq \phi(z) \quad \forall z \in M.$$

Then there is a unique minimizing geodesic  $\gamma : [0, 1] \rightarrow M$  between  $x$  and  $y$ . It is the projection of a normal extremal  $\psi : [0, 1] \rightarrow T^*M$  satisfying  $\psi(1) = (y, \frac{1}{2}D_y\phi)$ . In particular  $x = \exp_y(-\frac{1}{2}D_y\phi)$ .

# On the image of the exponential mapping

Proof.

Let  $y \neq x$  in  $M$  satisfying the assumption and  $\bar{\gamma} = \gamma_{\bar{u}} : [0, 1] \rightarrow M$  be a minimizing geodesic from  $x$  to  $y$ . We have for every  $u \in \mathcal{U} \subset L^2([0, 1]; \mathbb{R}^m)$  (close to  $\bar{u}$ ),

$$\|u\|_{L^2}^2 = C(u) \geq e_{SR}(x, E^{x,1}(u)) \geq \phi(E^{x,1}(u)),$$

with equality if  $u = \bar{u}$ . So  $\bar{u}$  is solution to the following **optimization problem**:

$\bar{u}$  minimizes  $C(u) - \phi(E^{x,1}(u))$  among all  $u \in \mathcal{U}$ .

We infer that there is  $p \neq 0$  such that

$$p \cdot D_u E^{x,1} = D_u C \quad \text{with } p = D_{E^{x,1}(u)} \phi$$

and in turn get the result. □

# On the image of the exponential mapping

## Remark

*If  $(M, d_{SR})$  is complete and there are no singular minimizing curves, then  $\exp_x(T_x^*M) = M$ .*

## Examples:

- Heisenberg.
- Fat distributions.
- For generic SR structures of rank  $\geq 3$ .

## Remark

*If  $(M, d_{SR})$  is complete and there are no strictly singular minimizing curves, then  $\exp_x(T_x^*M) = M$ .*

$\rightsquigarrow$  Medium fat distributions.

# Open problems in SR geometry I: The Sard conjecture

Let  $M$  be a smooth connected manifold of dimension  $n$  and  $\mathcal{F} = \{X^1, \dots, X^k\}$  be a family of smooth vector fields on  $M$  satisfying the Hörmander condition. Given  $x \in M$  and  $T > 0$ , the **End-Point mapping**  $E^{x,T}$  is defined as

$$\begin{aligned} E^{x,T} : L^2([0, T]; \mathbb{R}^m) &\longrightarrow M \\ u &\longmapsto x(T; x, u) \end{aligned}$$

where  $x(\cdot) = x(\cdot; x, u) : [0, T] \longrightarrow M$  is solution to the Cauchy problem

$$\dot{x} = \sum_{i=1}^m u_i X^i(x), \quad x(0) = x.$$

## Proposition

*The map  $E^{x,T}$  is smooth on its domain.*

# The Sard Conjecture

## Theorem (Morse 1939, Sard 1942)

Let  $f : \mathbb{R}^d \rightarrow \mathbb{R}^p$  be a function of class  $C^k$ , then

$$k \geq \max\{1, d - p + 1\} \implies \mathcal{L}^p(f(\text{Crit}(f))) = 0,$$

where  $\text{Crit}(f)$  is the set of critical points of  $f$ , i.e. the points where  $D_x f$  is not onto.

Let

$$\text{Sing}_{\mathcal{F}}^{x,T} := \left\{ u \in L^2([0, T]; \mathbb{R}^m) \mid u \text{ singular} \right\}.$$

## Conjecture

The set  $E^{x,T} \left( \text{Sing}_{\mathcal{F}}^{x,T} \right) \subset M$  has Lebesgue measure zero.

# Open problems in SR geometry II: Regularity of minimizing geodesics

Let  $(\Delta, g)$  be complete SR structure on a smooth manifold  $M$ .

## Open Question

*Do the minimizing geodesics enjoy some regularity ? Are they at least of class  $C^1$  ?*

↪ Very partial results by Monti, Leonardi and later Monti.

## References

- V. Jurdjevic. "Geometric Control Theory".
- A. Bellaïche. "The tangent space in sub-Riemannian geometry".
- R. Montgomery. "A tour of subriemannian geometries, their geodesics and applications".
- A. Agrachev, D. Barilari, U. Boscain. "Introduction to Riemannian and sub-Riemannian geometry".
- F. Jean. "Control of Nonholonomic Systems: From Sub-Riemannian Geometry to Motion Planning".
- L. Rifford. "Sub-Riemannian Geometry and Optimal Transport".

Thank you for your attention !!